



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

made of roots, moss and mud, well packed together, and owing to the daily showers, a nest generally raises a good crop of grass and weeds around its edges. The nest is placed commonly in coffee bushes, bunches of bananas or any low shrub at from eight to twelve feet from the ground. The extremes which I noted were two and twenty feet respectively. There were as many as 50 nests in the 80 acres surrounding the house. When the young are just beginning to fly the native small boy goes out and captures the little fellows by the dozen, using them to make bird stew. Gray's Robin lays two or three eggs, and although I have examined hundreds of nests I have never found four. There is great variation in size, shape and markings but an average egg resembles those of our Black-headed Grosbeak.

The Black-headed Saltator (*Saltator atriceps*) is a representative bird of Chiapas and is found mostly in flocks in the wooded portion of the lowlands and foothills, flitting about in the bushes and tree-tops chattering harshly all the time. The nest is loosely constructed of twigs and leaves and two eggs are always laid, being blue with heavy black markings at the larger end, similar to the eggs of the Redwinged Blackbird. The nesting season extends from the latter part of April to the latter part of July.

Synallaxis erythrorhax is a great deal like a little wren with nothing much to distinguish him but his nesting habits, but here he rivals anything in the feathered kingdom. The nest is usually placed from four to ten feet up in dense bushes and is composed of dry twigs four or five inches long. The bird makes a pile of these sticks a foot in diameter leaving a cavity of three or four inches diameter inside. After this is done a horizontal tunnel two inches in diameter is made, leading away from the nest. At about a foot from the nest at the end of the tunnel another pile of twigs is constructed through

which the tunnel turns upward for a few inches. This is the opening of the nest upon which the male roosts while the female is setting. The inner cavity is lined with leaves and three light blue eggs, rarely two, are laid.

I found the Grove-billed Ani (*Crotophaga sulcirostris*) rather common, nesting according to Davie as good birds should. I found sets varying from four to ten eggs and was told by Indians that 20 eggs were sometimes found in a nest, from which it appears that this species has the same habit as *C. ani* of several birds using the same nest. I found one nest of *C. sulcirostris* built on the top of a nest of Giraud's Flycatcher.

The Central American House Wren (*Troglodytes intermedius*) seemed like an old friend to me, flitting about and singing like our wrens at home. They reared their young in dark holes under the rafters of the house and barn and in the hollow trunks of trees. I found one nest in a hollow stump on the ground. The bird lays but three or four eggs, which are lighter than those of our common house wren. This little fellow and the Turkey Vulture were seemingly the only old friends I had in Mexico.



Scott's Oriole at San Diego, Cal.

SCOTT'S ORIOLE (*Icterus parisorum*) is not rare on the desert slope of the mountains in San Diego County in the migration, but it is very seldom seen between the mountains and the sea here. I saw two males in April, one of these being in the eucalyptus grove in the city park of San Diego. The song of this male was peculiar in some ways, so I followed him around to make sure of the identification. Last Sunday (June 2) I heard the same song in another part of the grove, a few hundred yards from where I saw the bird in April. This would indicate that he had become a summer resident here and probably had a mate. I know of no breeding record of this species in the coast region of San Diego County.

San Diego, Cal. FRANK STEPHENS.